

Treatment of Renal Colic in Emergency Hebron Government Hospital

Mohammad Qtait¹, Salah Tamiza²

¹Emergency Department, BSN, MSN in Hebron Government Hospital, Al Quds University, Health Professional Collage, Hebron, Palestine

²Emergency Department, Emergency Specialist in Hebron Government Hospital, Hebron, Palestine

Email address:

mohamadtaha98@hotmail.com (M. Qtait)

To cite this article:

Mohammad Qtait, Salah Tamiza. Treatment of Renal Colic in Emergency Hebron Government Hospital. *International Journal of Clinical Urology*. Vol. 1, No. 1, 2017, pp. 11-14. doi: 10.11648/j.ijcu.20170101.13

Received: May 19, 2017; **Accepted:** May 31, 2017; **Published:** July 18, 2017

Abstract: The aims the present investigation aimed to point out some data concerning renal colic in Palestine, and to evaluate the practical clinic approach to it. All the renal colic cases treated in three month in the Hebron hospital ED, reviewed to assess the data, diagnostic and treatment patterns, and to compare them with what is reported in the literature. Retrospective study by review of file of emergency department in three month of renal colic or urinary stone. Renal colic were diagnosed in three month 300 cases (1%) from all vister of ED; 66% of them were recurrent stone formers; the males 61%, female 39%, The age distribution, showed a higher rate from 25 to 44 years of age, Ultrasonography (US) was the examination in 72% cases, 99% lab test urine analysis. Diclofen were always used (90%), association with Hyoscine butylbromide, narcotic as pethidin (10%). The data of our investigation are in a substantial agreement with the reported literature bout features of renal colic and its treatment. The diagnostic approach is mainly based on US and lab test. CT-scan was not used in ED. Conclusion; the data of our investigation agree substantially with those reported in the literature as far as concerns of renal colic and its treatment. The diagnostic approach is mainly based on US whereas urine examination, most of patient given Diclofen, and Hyoscine butylbromide and 10% given narcotic. In pale stain treated the renal colic as national guide.

Keywords: Renal Colic, Stones, Emergency Department

1. Introduction

Pain is a most reason for coming to emergency departments (EDs) [12]. Stone in urinary system is a common disease in emergency department ED with prevalence between 7 – 21 cases per 10.000 in habitants in Western countries [1]. "Renal colic is defined as severe intermittent flank pain that radiates to the groin, lower abdomen, or genitalia due to the passage of a stone through the urinary system. The signs characterized by acute flank pain which sometimes fluctuates an radiates to the groin and scrotum in males" [15]. The stones form different causes; genetic, metabolic, socio-economic, cultural and geographic factors [3]

Hebron is the largest city in the West Bank and have a population of around 706.000 (PBCS report, 2015). These people are the target centers of the service offered by the hospitals operating in Hebron. In Hebron, there are governmental and non-government hospitals that operates at

high capacity to meet the population demands. Hospitals in Hebron operate at the limits of their effective capacity or even they operate over capacity in some of the hospital departments. Alia Hospital is the most crowded in Palestine, operating at 124%, average occupancy rate in 2015 [11] The second hospital in the West bank crowded ED is Hebron hospital is the large in 2015. 60750 patient seen in ED, More than 1000 patient complain from renal colic [11]. No papers have been published treatment of renal colic in Palestine emergency departments.

Renal colic pain is a described as severe pain the patient has ever experienced, Consequently, the use of effective pain killers like; non-steroidal anti-inflammatory drugs (NSAIDS), and narcotic, or a combination of medications (anti-inflammatory and spasmolytic agents), play important roles in the treatment of these patients [8].

The effect of NSAIDs on relieving pain in renal colic is similar to opiates, the disadvantage of NSAIDs, in the oral or rectal form, is the delayed onset time compared with

intravenous morphine. Intravenous forms of NSAIDs are available and have a rapid onset, the side effects from the intravenous form of NSAIDs have been reported more frequently than for other types of drugs, complications of NSAIDs include; nausea, vomiting, feeling of heat or pressure in the chest, fatigue and lethargy [3].

NSAIDs have the advantage of decreasing ureteral smooth muscle tone, thereby In these cases other analgesics should be used. Diclofenac is a common NSAID used in patients with renal colic.

Both parenteral opioids and non-steroidal anti-inflammatory drugs (NSAIDs) are commonly used to provide relief from renal colic, and both can have adverse effects

Although opioids are effective and provide rapid analgesia in renal colic, hypotension, nausea, vomiting and dizziness are not uncommon after narcotic administration. Paracetamol is analgesics are commonly employed in combination with spasmolytic drugs for the treatment of renal and ureteral colic.

The involvement of prostaglandins promoted clinical experimentation to investigate the effectiveness of non-steroidal anti-inflammatory drugs (NSAID) in alleviating the pain of renal and ureteral colic.

Although the efficacy of spasmolytic agents in renal and ureteral colic has been debated, they are widely used alone or in combination with narcotic analgesics in many countries including Saudi Arabia and other Arabian Gulf countries. In a renal colic study, a single 20-mg intravenous dose of hyoscine N-butylbromide (Buscopan) demonstrated that this spasmolytic agent has a low analgesic effect when it is used as single therapy or with a 2.5-gm dose of dipyron [9].

Hyoscine Butyl Bromide is anti-muscarinic agents are effective in the treatment of smooth muscle spasms (especially gastrointestinal). Ureteral peristaltic activity of the genitourinary system is controlled by the autonomic nervous system so the use of anti-muscarinic agents can be effective. [5]

The present investigation aimed to point out some data concerning renal colic in Palestine, and to evaluate the practical clinic approach to it. For this reason all the renal colic cases treated in three month in the Hebron hospital ED have been reviewed to assess the data, diagnostic and treatment patterns, and to compare them with what is reported in the literature.

2. Purpose of the Study

The purpose of study present investigation aimed to point out some data concerning renal colic in Palestine, and to evaluate the practical clinic approach to it.

3. Method

3.1. Study Design

Retrospective study, take the medical records of the ED in Hebron Hospital, from 1 may to 31 July 2016, were

reviewed. All the records with a final diagnosis of definite renal colic or urinary tract stone were selected and collected for the analysis. The records were carefully reviewed and unconfirmed diagnoses such as, for example: “suspected renal colic” or “pain probably due to urinary tract disease”, were excluded from the final analysis.

3.2. Study Period

The date and time of visit was recorded together with the patient’s age and sex, and, if any, previous history of renal colic or urinary stones. Information concerning the clinical presentation, namely abdominal or lumbar pain, side of the pain, association with hematuria or accompanying symptoms such as nausea, vomiting or fever, were recorded as well. In The diagnostic procedures used were specified in every patients, namely US, and urine examination. Similarly, the given therapy, namely diclofen, antispastic and analgesic medications, opioids, were recorded. Lastly, the outcome was reported, that is admission, discharge or lost at follow up.

3.3. Inclusion Criteria

Were at least 18 years of age (and less than 65 years), Exclusion criteria were: not diagnosis renal stone, renal failure, diabetes, hypertension, pregnant and breast feeding women.

3.4. Study Setting

This study was conducted in the West bank in emergency department Hebron government hospital south of Palestine.

3.5. Ethical Consideration and Accessibility

Permission obtained to access the MOH hospitals report when approval by the director of hospital services. Not use the name or taken the name of participant.

3.6. Data Analysis Procedure

After data collection, from file filled in the table and entered and analyzed using the Statistical Package for Social Science program (computer software SPSS V. 22) for descriptive and inferential statistics. Frequencies were used to present the distribution of study variables.

4. Result

There were a total of 7,623 ED visits recorded from 1 May to 31 July 2016. Renal colic episodes were diagnosed in 300 cases (1%); 198 (66.6%) had a previous history of renal calculi and 183 (61%) were men, 39% female.

The age distribution shows a higher rate among young adults (from 25 to 45 years of age). The seasonal distribution outlines the maximum risk during summer months, particularly in July, and the minimum number of cases during the winter months.

Pain accounted for the chief complaint in all the cases, and was associated with nausea and vomiting in 39% of cases.

Fever was present in five patients (1%). Gross hematuria was observed in only 4% of cases, but microscopic hematuria was present in 92% of the urine performed. Total of 99% had urine analysis, and CBC, 69% had BUN and creatinine, US was the sole examination in 72% of cases.

About 12% of the patients did not receive any drugs in the ED either because they were already treated at home or any out medical center, or because pain had subsided in the ED.

300 patients (96%) were discharged, 12 patients (4%) were admitted to the urology ward after an urologic consultation (mostly for refractory pain, fever or ultrasound evidence of severe obstruction).

Table 1. Demographics.

Items	Percent	frequency
Gender		
Male	61%	183
Female	39%	117
Age		
less than 30	44.4%	133
Between 30-44	43.3%	130
More than 45	12.2%	37
colic		
First	33.4%	
Recurrent	66.6%	
Laterality of stone		
Left side	59%	
Right side	41%	

Table 1. Continue.

Shift	Number	Percent
Day	96	32%
Evening	132	44%
Night	72	24%

Table 2. Method of diagnosis.

Method of diagnosis	Lab test	Radiology US	Other
Urine analysis	99	72	2

Table 3. Type of medication use.

Drug use	Percent	No
Hyoscine butylbromide,	99%	
Pethidin,	10%	
Diclofen sodium,	90%	
Take all	10%	

When therapy was given, NSAIDs (diclofen sodium) were always used (90%) in association with opiate (tramadol) (9%) and 96% from patient given (hyoscine butylbromide). 10% given pethidin, 10% given all (diclofen sodium, hyoscine butylbromide, pethidin).

5. Discussion

One of the primary goals of emergency department staff is the prompt, and remove suffering from pain for patient coming in ED. The goal have been poor both in provision of analgesia and delay to analgesia [6]. The results of our

investigation confirm that renal colic pain is a quite frequent cause for ED visits. Since the Hebron Hospital inhabitants area, an annual incidence of 1% inhabitants may be estimated. Therefore, the epidemiological features we observed in our region are very similar to that reported in the literature [3].

Diagnosed in 300 cases (1%) from total this rate of renal colic or stone (1%) is the same that was reported in world study involving more than 100 million emergency department patient presentations [2].

Even if in many cases the clinical presentation was sufficient to diagnose a renal colic episode, US was performed in 72% of our patients due to less cost for hospital and no side effect on patient. Instead, a KUB was not performed and not request ED of them. This is not in keeping with Italian guidelines that suggest US as the initial diagnostic steps because the combination increases the sensitivity and specificity for urinary tract stone diagnosis [4]. On the other hand, US is a quick, simple, accurate and low cost procedure.

90% from patient given NSAIDs, 99% from patient use Hyoscine butylbromide, less cost and effect and not effect in balance as narcotic and that accept with study to Regarding treatment for renal colic, our data are in keeping with the recommendations of the literature. Several studies show that NSAIDs administration is the first choice of analgesia for acute renal colic, and they are at least as effective as narcotics [5]

The urinalysis is use for every patient complain from abdominal pain and its available in an emergency setting and doctors request its not cost and see the microscopic hematuria before US because of the need to have filled bladder for this examination.

This study does not seen report use of calcium channel blockers or a-antagonists, and rom study the physician use the same protocol in the treatment of renal colic.

The most of patients were able to be discharged. Admission was only required for complicated cases (fever, single kidney obstruction) or for those patients who could not achieve pain relief.

In conclusion, the data of our investigation are in a substantial agreement with those reported in the literature as far as concerns of renal colic and its treatment. The diagnostic approach in Hebron hospitals is mainly based on US whereas urine examination, most of patient given NSAIDs, and Hyoscine butylbromide and 10% given narcotic. In pale stain treated the renal colic as national guide. The most use diagnosis laboratory test urine analysis then US in ED.

6. Conclusion

Renal colic were diagnosed in three month 300 cases (1%) from all vister of ED; 66% of them were recurrent stone formers; the males 61%, female 39%, The age distribution, showed a higher rate from 25 to 44 years of age, Ultrasonography (US) was the examination in 72% cases,

99% lab test urine analysis. Diclofen were always used (90%), association with Hyoscine butylbromide, narcotic as pethidin (10%). The data of our investigation are in a substantial agreement with the reported literature about features of renal colic and its treatment. The diagnostic approach is mainly based on US and lab test. CT-scan was not used in ED. The data of our investigation are in a substantial agreement with those reported in the literature as far as concerns of renal colic and its treatment. The diagnostic approach is mainly based on US whereas urine examination, most of patient given Diclofen, and Hyoscine butylbromide and 10% given narcotic. In pale stain treated the renal colic as national guide.

Abbreviation

MOH	Ministry of health
PBCS	Palestinian Central Bureau of Statistics
ANOVA	Analysis of Variances.
ED	emergency department
NSAIDS	non-steroidal anti-inflammatory drugs
US	Ultra sound

References

- [1] Amato M, Lusini ML, Nelli F (2004) Epidemiology of nephrolithiasis today. *Urol Int* 1: 1–5.
- [2] Brown J (2006) Diagnostic and treatment patterns for renal colic in US Emergency Departments. *Int Urol Nephrol* 38: 87–92.
- [3] Coe FL, Parks JH, Asplin JR (1992) The pathogenesis and treatment of kidney stones. *N Engl J Med* 327: 1141–1152.
- [4] Davenport K, Timoney AG, Keeley FX (2005). Conventional and alternative methods for providing analgesia in renal colic. *BJU Int.*; 95(3): 297–300.
- [5] Davenport K, Waive E. The Role of Non-Steroidal Anti-Inflammatory Drugs in Renal Colic. *Pharmaceuticals*. 2010; 3(5): 1304–1310.
- [6] Ducharme J, Barber C. (1995) A prospective blinded study of emergency pain assessment and.
- [7] Ducharme J. (1994) Emergency pain management: A Canadian Association of Emergency.
- [8] Holdgate A, Oh CM. Is there a role for antimuscarinics in renal colic? A randomized controlled trial. *J Urol*. 2005; 174(2): 572–5.
- [9] Lloret J, Munoz J, Monmany J, Puig X, Bonastre M, Brau J, Sola J, Domingo P, Jane F. Treatment of renal colic with dipyrone. *Curr Ther Res* 1987; 42: 1119-26.
- [10] Lundstam S, Wahlander L, Leissner KH, Karl JG. Prostaglandin-synthetase inhibition with diclofenac sodium in treatment of renal colic: comparison with use of a narcotic analgesic. *Lancet* 1982; 1: 1096-7.
- [11] MOH, annual report, 2015.
- [12] Paris PM, Stewart RD (1988) (Eds.) Pain management in emergency medicine. Norwalk, CT. Appelton and Lange. Physicians consensus document *Journal of Emergency Medicine* 12: 855–66.
- [13] Samad EJ Golzari, Hassan Soleimanpour, Farzad Rahmani, Nahid Zamani Mehr, Saeid Safari, Yaghoob Heshmat, Hanieh Ebrahimi Bakhtavar 2014 Feb; Therapeutic Approaches for Renal Colic in the Emergency Department: A Review Article4(1): e16222. Published online 2014 Feb 13. doi: 10.5812/aapm.16222 PMID: PMC3961032.
- [14] Sanahuja J, Corbera G, Garau J, Pl R, Carmen Carre M. Intramuscular diclofenac sodium versus intravenous baralgin in the treatment of renal colic. *DICP, The Annals of Pharmacotherapy* 1990; 24: 361-4.
- [15] Seifter JL, Brenner BM. Urinary Tract Obstruction. *Harrison principles of internal medicine*; 17th ed. Braunwald: McGraw Hill, 2008; 1827-30.
- [16] Serinken M, Karcioğlu O, Turkcuer I, Ozkan HI, Keysan MK, Bukiran (2008) A. Analysis of clinical and demographic characteristics of patients presenting with renal colic in the emergency department. *BMC Res Notes.*; 1: 79. doi: 10.1186/1756-0500-1-79 therapy. *Journal of Emergency Medicine* 13: 571.
- [17] Thompson JF, Pike JM, Chumas PD, Rundle JSH. Rectal diclofenac compared with pethidine injection in acute renal colic. *Br Med J* 1989; 299: 1140-1.